

Bexidal®

Tablet

Description

Bexidal is a preparation of Mebhydrolin as Mebhydrolin Napadisylate BPC.

Indications

Allergic diseases or symptoms, such as urticaria, pruritus of various origin, eczema, drug rash, allergic conjunctivitis, dermatitis of nutritional origin, hay fever, vasomotor rhinitis, allergic asthma.

Dosage and Administration

Unless otherwise prescribed, the following doses are recommended :

Adults and children over 10 years	2-6 tablets daily
Children from 5-10 years	2-4 tablets daily
Children from 2-5 years	1-3 tablets daily
Children up to 2 years	1-2 tablets daily

Treatment should be given in several single doses daily. Bexidal may be swallowed during or shortly after meals. For children, the tablets may be crushed and mixed with food.

Bexidal is essentially free from secondary sedative-hypnotic effects. It does not impair psycho-physical efficiency to any appreciable degree.

Contraindications

- ◆ Caution is required while driving or operating heavy machinery
- ◆ Concomitant use of CNS depressant or alcohol
- ◆ Patients to whom Mebhydrolin has previously been suspected to have agranulocytosis or neutropenia
- ◆ First trimester of pregnancy

Commercial Pack

Bexidal® Tablet : Box containing 20 aluminium strips of 10 tablets. Each tablet contains Mebhydrolin 50 mg as Mebhydrolin Napadisylate BPC.