

Emonium[®]

Tiemonium Methylsulfate
Tablet

Description

It is an antispasmodic. It reduces muscle spasms of the intestine, biliary system, bladder and uterus.

Indications

This drug is indicated for pain in gastrointestinal and biliary diseases, and in urology and gynaecology.

Dosage and Administration

In adults the usual dose is 2 to 6 tablets per day.

Contraindications

This drug should not be used in glaucoma (acute pain of eyeball with vision disturbances) and in difficulty to urinate (disorders of prostate or bladder).

Precautions

In order to avoid possible drug interactions, you must systematically inform your physician about any other drugs you are taking.

Use in Pregnancy and Lactation

Avoid taking this drug without medical advice during pregnancy or breast feeding.

Adverse Effects

As with any active substance this drug may cause certain undesirable or unpleasant effects. In some people dry mouth with difficulty in swallowing and thirst, dilatation of pupils with loss of accommodation and sensitivity to light, increased intraocular pressure, flushing,

dry skin, bradycardia followed by tachycardia, palpitations and arrhythmias, difficulty in micturition and constipation may occur.

Overdose

Occasional retention of urine in excessive or overdoses.

Pharmaceutical Precautions

Keep in a cool, dry place in accordance with good pharmaceutical practice. Keep out of the reach of children.

Commercial Pack

Emonium[®] tablet : Box containing 50 tablets in 5 x 10's blister strips. Each tablet contains Tiemonium Methylsulfate INN 50 mg.



Manufactured by

BEXIMCO PHARMACEUTICALS LTD.

TONGI, BANGLADESH

BL 4547

300806

© Emonium is a registered trademark of Beximco Pharmaceuticals Ltd.