

Metazine® MR

Trimetazidine Hydrochloride
Tablet

Description

Trimetazidine Hydrochloride is a unique antianginal and anti-ischemic agent that exerts anti-ischemic effects without inducing hemodynamic changes and improves the status of ischemic myocardium. The drug exerts its antianginal effects through a direct cytoprotective action on the myocardium, thereby avoiding the side effects of impaired left ventricular function, excessive vasodilation and other disadvantages associated with the use of comparator antianginal agents.

Indications

Trimetazidine Hydrochloride is indicated for-

1. Ischemic heart disease (angina pectoris, sequelae of infarction)
2. Adjuvant symptomatic treatment of vertigo and tinnitus
3. Adjuvant treatment of the decline visual acuity and visual field disturbances, presumably of vascular origin.

Dosage And Administrations

One tablet at mealtimes in the morning and evening.

Side effects

Trimetazidine Hydrochloride is safe and well tolerated. The most common side effects encountered are gastric discomfort, nausea, headache and vertigo. However, the side effects are mild and non-specific.

Contraindications

Trimetazidine Hydrochloride is contraindicated in patients who are hypersensitive to Trimetazidine.

High Risk Group

Use in Pregnancy:

Studies in animals have not demonstrated teratogenic effects. However, in the absence

of clinical data and for safety reasons, prescription should be avoided during pregnancy.

Use in Nursing mother:

There is no information on the secretion of Trimetazidine Hydrochloride into breast milk. However, breast feeding should be discontinued if the use of Trimetazidine Hydrochloride is considered essential.

Use in Children:

Safety and efficacy have not been established with Trimetazidine Hydrochloride use in children.

Drug Interactions

No drug interactions so far have been reported. In particular, no interactions have been reported with beta-blockers, calcium antagonists, nitrates, heparin, hypolipidemic agents or digitalis preparation.

Pharmaceutical Precautions

Keep in a cool & dry place. Protect from light. Keep out of the reach of children.

Commercial Pack

Metazine® MR Tablet: Box containing 30 tablets in 3X10's Alu-Alu form packs. Each modified release tablet contains Trimetazidine Hydrochloride BP 35 mg.



Manufactured by

BEXIMCO PHARMACEUTICALS LTD.

TONGI, BANGLADESH

BL 8342

271112

© Metazine is a registered trademark of Beximco Pharmaceuticals Ltd.

