

Tenafin[®]

Butenafine Hydrochloride
Cream

Description

Butenafine Hydrochloride is a broad-spectrum antifungal agent. Butenafine Hydrochloride is hypothesized to act by inhibiting the epoxidation of squalene, thus blocking the biosynthesis of ergosterol, an essential component of fungal cell membranes.

Indication

Tenafin[®] (Butenafine Hydrochloride) cream is indicated for the topical treatment of the following superficial dermatophytosis:

- interdigital tinea pedis (athlete's foot),
- tinea corporis (ringworm) and
- tinea cruris (jock itch) due to *E. floccosum*, *T. mentagrophytes*, *T. rubrum*, and *T. tonsurans*.

Dosage and Application

In the treatment of interdigital tinea pedis, Tenafin[®] should be applied twice daily for 7 days or once daily for 4 weeks. Patients with tinea corporis or tinea cruris should apply Tenafin[®] once daily for two weeks.

Sufficient Tenafin[®] cream should be applied to cover affected areas and immediately surrounding skin of patients with interdigital tinea pedis, tinea corporis and tinea cruris.

Contraindications

Tenafin[®] (Butenafine Hydrochloride) is contraindicated in individuals who have known or suspected sensitivity to Tenafin[®] cream or any of its components.

Warnings and Precaution

Tenafin[®] (Butenafine Hydrochloride) is not for ophthalmic, oral, or intravaginal use. Tenafin[®] cream is for external use only. If irritation or

sensitivity develops with the use of Tenafin[®] cream, treatment should be discontinued and appropriate therapy instituted.

Adverse Reactions

Rarely local mild burning or irritation may be experienced. Hypersensitivity reactions may occur.

Pharmaceutical Precautions

Store between 5° and 30°C.

Commercial Pack

Tenafin[®] cream : Tube containing 15 g cream, each gram contains Butenafine Hydrochloride INN 10 mg.



Manufactured by

BEXIMCO PHARMACEUTICALS LTD.

TONGI, BANGLADESH

BL 4181

111006

® Tenafin is a registered trademark of Beximco Pharmaceuticals Ltd.